


Forward Physics in Herwig++

Pavel Ruzicka

Institute of Physics, AS CR


Forward Physics at the LHC, Manchester
December 13, 2010



Introduction

Herwig++ has been being extended for some of the most interesting forward physics processes:


- Two photon physics
- Hard inclusive diffraction: Single Diffraction, Double Pomeron Exchange
- Central Exclusive Production



ThePEG structure

- Herwig++ is based on thePEG framework: the Toolkit for High Energy Physics Event Generation.
- ThePEG provides all the infrastructure that does not depend on the physics models. Each HERWIG++ class inherits from ThePEG abstract class and provides specific model implementation.
- ThePEG has repository which keeps all the building blocks of the EventGenerator and connections between them (as Matrix elements, PDF functions, hadronization handlers, beam properties etc.).
- User setups all the building blocks and their properties easily together via the text inputfile.

Two photon physics



Budnev flux:

$$dN = \frac{\alpha}{\pi} \frac{dE_\gamma}{E_\gamma} \frac{dQ^2}{Q^2} \left[\left(1 - \frac{E_{\text{gamma}}}{E}\right) \left(1 - \frac{Q^2}{Q_0^2}\right) F_E + \frac{E_\gamma^2}{E^2} F_M \right]$$

$$F_M = G_M^2$$

$$F_E = (4m_p^2 G_E^2 + Q^2 G_M^2) / (4m_p^2 + Q^2)$$





$$G_E^2 = G_M^2 / \mu_p^2 = (1 + Q^2 / Q_0^2)$$

where $\mu_p^2 = 7.78$ and $Q_0^2 = 0.71 \text{ GeV}^2$

Two photon physics

- Budnev flux implemented.
- Available SM processes: dilepton production, W boson production, dijet production
- Extension for beyond standard model processes can be easily done.
- Two photon physics validated against FPMC (Forward Physics Monte Carlo, www.cern.ch/fPMC, arXiv:0903.3861), see next slides and LPAIR.

Distributions $pp \rightarrow p + \gamma\gamma + p \rightarrow p + \mu\mu + p$




Cross Section $pp \rightarrow p + \gamma\gamma + p \rightarrow p + \mu\mu + p$

p_T	Herwig++	FPMC
3 GeV	47.2 pb	48.9 pb
5 GeV	13.2 pb	13.5 pb
10 GeV	2.20 pb	2.22 pb
30 GeV	1.09×10^{-2} pb	1.09×10^{-2} pb
50 GeV	2.48×10^{-3} pb	2.49×10^{-3} pb

Table: Cross sections of dimuon production for various p_T cuts at $\sqrt{s} = 14$ TeV with $x_{min} = 0$, $x_{max} = 1$ and $Q^2 = 2$ GeV²

Inclusive diffraction: Ingelman-Schlein model



- Cross section:

$$\sigma(P\bar{P} \rightarrow P\bar{P}XY) = \sum_{i,j} \iint dx_1 dx_2 f_i^D(x_1, Q_1^2) f_j^D(x_2, Q_2^2) \hat{\sigma}(x_1, Q_1^2, x_2, Q_2^2)$$





$$f_i^D(x, Q^2) = f_{\mathbb{P}}(t, x_{\mathbb{P}}) f_i^{\mathbb{P}}(Q^2, x) + f_{\mathbb{R}}(t, x_{\mathbb{R}}) f_i^{\mathbb{R}}(Q^2, x)$$

- Pomeron/Reggeon flux: $f_{\mathbb{P}}(t, x_{\mathbb{P}}) = A e^{\beta_{\mathbb{P}} t} / x^{2(\alpha(0) - \alpha' t) - 1}$
- Pomeron/Reggeon structure function: $f_i^{\mathbb{P}}(Q^2, x)$





Current implementation

- Sum of pomeron and reggeon contributions implemented.
- Default value of pomeron and reggeon fluxes are given by the PDF fits. They can be also varied by user.
- Pomeron structure functions from HERA measurement: 2006 (fit A and fit B), 2007.
- The PDFs can be freezed or extrapolated outside of their boundaries.
- In case of reggeon, pion structure function is used.
- Pomeron composed either from the valence gluons (default option) or from $q\bar{q}$ pairs.
- Reggeon has only $q\bar{q}$ structure which is consistent with pion.
- Gap survival probability is not included.
- Comparison was done with POMWIG (the most established event generator for hard diffraction, <http://www.pomwig.com>).

Distributions $pp \rightarrow p + \text{dijet} + p$



Distributions $pp \rightarrow p + dijet + p$




Cross Section $pp \rightarrow p + dijet + p$

p_T	Herwig++	Pomwig
10 GeV	9.63×10^4 nb	9.56×10^4 nb
20 GeV	7.49×10^3 nb	7.49×10^3 nb
30 GeV	1.55×10^3 nb	1.54×10^3 nb
40 GeV	2.83×10^2 nb	2.81×10^2 nb

Table: Cross sections of dijet production for various p_T cuts at $\sqrt{s} = 14$ TeV with $x_{min} = 10^{-7}$, $x_{max} = 1$ H1 2006 Fit A.


Central Exclusive Production



- KMR model for Central Exclusive Production.
- Processes plan to be added: Higgs, dijets, di-photons, $q\bar{q}g$, $Wq\bar{q}$
- Validated against ExHuME





$$M^2 \frac{\delta\sigma}{\delta t_1 \delta t_2 \delta y \delta M^2} = M^2 \frac{\delta\mathcal{L}}{\delta t_1 \delta t_2 \delta y \delta M^2} \hat{\sigma}$$

KMR luminosity



$$M^2 \frac{\delta \mathcal{L}}{\delta t_1 \delta t_2 \delta y \delta M^2} = S e^{b(t_1+t_2)} \left[\frac{\pi}{8} \int \frac{dQ_t^2}{Q_t^4} f_g(x_1, Q_t^2) f_g(x_2, Q_t^2) \right]$$

Higgs production



Summary & Outlook

Summary:

- Two photon physics is available: 2 fermion \rightarrow 2 fermion processes, W production available.
- Hard inclusive diffraction added: SD, DPE: dijet production, W and Z production, Higgs production.
- All tested against: POMWIG, FPMC and LPAIR
- KMR luminosity implemented: Higgs production (not in release yet)

Outlook:

- Add more processes into CEP: dijet production, di-photon, $q\bar{q}g$, $Wq\bar{q}$ (can be expected in first half of next year)