

Low-energy Lepton Observables in the MRSSM

Hyejung Stöckinger-Kim

collaboration with D. Stöckinger and W. Kotlarski

arXiv:1902.0665

TU Dresden

PASCOS2019
Manchester, 1. July 2019

Low-energy lepton observables

$$(g-2)_\mu: a_\mu^{\text{Exp. BNL(E821)}} - a_\mu^{\text{SM}} = (28.1 \pm 6.3^{\text{Exp.}} \pm 3.6^{\text{Th(KNT18)}}) \times 10^{-10}$$

Keshavarzi, Nomura, Teubner'18; Jegerlehner'17: $\pm 4.4^{\text{Th}}$

New experiment at **Fermilab(E969)**: ~ 0.14 ppm

$$\mu \rightarrow e\gamma: B_{\mu^+ \rightarrow e^+ \gamma} = \frac{\Gamma(\mu \rightarrow e\gamma)}{\Gamma(\mu \rightarrow e\nu_\mu \bar{\nu}_e)} < 4.2 \times 10^{-13} \text{ MEG 2016}$$

Future **MEG-II**: expect to improve sensitivity $\sim 5 \times 10^{-14}$

$$\mu \rightarrow e: B_{\mu\text{Au} \rightarrow e\text{Au}} < 7 \times 10^{-13} \text{ SINDRUM 2006}$$

Future **COMET** and **Mu2E**: expected 7.2×10^{-15}

- Parameter ranges? Correlation among them?
- LFV processes $\propto \delta_{12}^{\text{L/R}}$ (1. and 2. generation slepton mixing)

Outline

- 1 Model: MRSSM
- 2 $g - 2$ and $\mu \rightarrow e\gamma$
- 3 $\mu \rightarrow e$
- 4 Correlation between the observables
- 5 Summary

Minimal R-Symmetric Supersym. SM

MSSM with a continuous unbroken R-symmetry $U(1)_R$: R-charge

[Kribs, Poppitz, Weiner]

$$\text{enlarged } \begin{cases} \text{Higgs sector: } \hat{R}_{u,d} \\ \text{Gauge sector: } \hat{O}, \hat{T}, \hat{S} \end{cases}$$

Fields	Fermion		Boson	
Matter	l, e_R^*	0	\tilde{l}, \tilde{e}_R^*	+1
	q, d_R^*, u_R^*	0	$\tilde{q}, \tilde{d}_R^*, \tilde{u}_R^*$	+1
Gauge	$\tilde{g}, \tilde{W}, \tilde{B}$	+1	g, W, B	0
Adjoint chiral	$\tilde{O}, \tilde{T}, \tilde{S}$	-1	O, T, S	0
H-Higgs	$\tilde{H}_{u,d}$	-1	$H_{u,d}$	0
R-Higgs	$\tilde{R}_{u,d}$	+1	$R_{u,d}$	+2

$$\tilde{e}_R, \tilde{u}_R, \tilde{d}_R : -1$$


MRSSM Superpotential and Interaction terms

$W_{\text{sup.pot.}}^{\text{MRSSM}} \ni \Lambda_d \hat{R}_d \cdot \hat{T} \hat{H}_d + \lambda_d \hat{S} \hat{R}_d \cdot \hat{H}_d + (d \rightarrow u)$ Yukawa-like sup.pot. parameters:

$$\mathcal{L}_{\text{int}} \ni \sum_{(f, \tilde{f}), g} \left(\overline{\tilde{\chi}}_A^0 [n_{gA}^{\text{L}(f)} \text{P}_L + n_{gA}^{\text{R}(f)} \text{P}_R] f_g \tilde{f}_R^\dagger + \overline{\tilde{\chi}}_A^{0c} [o_{gA}^{\text{L}(f)} \text{P}_L + o_{gA}^{\text{R}(f)} \text{P}_R] f_g \tilde{f}_L^\dagger \right) \\ + \sum_{(f', \tilde{f}'), g} \left(\overline{\tilde{\chi}}_A^- [c_{gA}^{\text{L}(f')} \text{P}_L + c_{gA}^{\text{R}(f')} \text{P}_R] f'_g \tilde{f}'_L^\dagger \right)$$


Gauginos interactions: $n^{\text{R}}, o^{\text{L}}, c^{\text{L}}$

Higgsino interactions: $n^{\text{L}}, o^{\text{R}}, c^{\text{R}} \leftarrow$ suppressed by small Yukawa couplings



MRSSM: Λ_d, λ_d , no $\tan \beta$ -enhancement like MSSM .

Observables and diagrams



$$A_1^{ji} [q^2 \gamma^\mu]$$

$$A_Z$$

$$A_{\text{box}}$$

$$A_2^{ji} [\sigma^{\mu\nu} q_\nu]$$

$$A_1^{\bar{e}\mu}, A_2^{\bar{e}\mu}$$

$$(g-2)_\mu \propto A_2^{\bar{\mu}\mu}$$

$$\mu \rightarrow e\gamma \propto A_2^{\bar{e}\mu}$$

$$\mu \rightarrow e \propto A_1^{\bar{e}\mu}, A_2^{\bar{e}\mu}, A_Z, A_{\text{box}}$$

- 1 a_μ and $g \rightarrow e\gamma$ always correlated
- 2 correlation of a_μ and $\mu \rightarrow e$
 \uparrow when dipole A_2 dominates
- 3 large λ_d or Λ_d
- 4 $\mu \rightarrow e\gamma, \mu \rightarrow e \propto \delta_{12}^{L,R}$
 slepton-generation-mixing

Survey of relevant parameters

$$M_B^D, M_W^D, \mu_d, \lambda_d, \Lambda_d, m_{\tilde{l},22}, m_{\tilde{e},22} \quad ((m_{\tilde{e}})_{11} = (1.5)m_{\tilde{e},22}, (m_{\tilde{l}})_{11} = (1.5)m_{\tilde{l},22})$$

dimensionless LFV parameters: $\delta_{12}^L \equiv \frac{(m_{\tilde{l}}^2)_{12}}{m_{\tilde{l},11}m_{\tilde{l},22}}, \delta_{12}^R \equiv \frac{(m_{\tilde{e}}^2)_{12}}{m_{\tilde{e},11}m_{\tilde{e},22}}$






Scenario	light masses			
BHL	$M_B^D, \mu_d, m_{\tilde{l}}$	$\delta_{12}^R \neq 0$	WHR, BHR	BR
BHR	$M_B^D, \mu_d, m_{\tilde{e}}$	$\delta_{12}^L \neq 0$	BHL	BL, WL
WHL	$M_W^D, \mu_d, m_{\tilde{l}}$			
Equal-mass	$M_B^D, M_W^D, \mu_d, m_{\tilde{l}}, m_{\tilde{e}}$			

λ 's, Λ 's: perturbative ($|\lambda_i|, |\Lambda_i| < 4$)

$|\lambda_u|, |\Lambda_u| < 2$ from EW prec. T-parameter

LHC: small mass splitting between sparticles, $M_{\tilde{\chi}_1^\pm} > 92$ Gev, $m_{\tilde{\mu}_R} > 94$ Gev

Numerical analysis of $g - 2$





Smaller hyper-charge: BHL < BHR
 WHL ← larger SU(2) gauge coupling
 λ_d , Λ_d enhancement: WHR, BHR, BHL
 BL, BR, WL: simple, mainly gauge couplings

Equal mass cases
 upper 3 bands: $\Lambda_d = 4$
 lower 3 bands: $\Lambda_d = 0$

a_μ and $\mu \rightarrow e\gamma$

$$a_\mu \propto A_2^{\bar{\mu}\mu L} + A_2^{\bar{\mu}\mu R}$$

$$B_{\mu \rightarrow e\gamma} \propto |A_{2\text{red}}^{\bar{e}\mu L}|^2 \times |\delta_{12}^L|^2 + |A_{2\text{red}}^{\bar{e}\mu R}|^2 \times |\delta_{12}^R|^2$$






$\mu \rightarrow e$ with respect to $\tan \beta$, $m_{\tilde{q}}$, λ_u

Useful quantity: $R(N) \equiv \frac{B_{\mu N \rightarrow e N}}{B_{\mu \rightarrow e \gamma}}$; dependence on the δ 's drops out

Especially for A_2 dominance: dipole A_2 form factor drops out; perfect correlation; only dependent on the element used in the experiment. [Kitano, Koike, Okada, 2002]

$$R^{\text{only dip.}}(Al) = 0.0026.$$




→ MEG limit on $\mu \rightarrow e \gamma$ determines the maximum possible $\mu \rightarrow e$ conversion rate

Impact of the additional form factors A_1 , A_Z , A_{box} → deviation

Large $\lambda_u \rightarrow A_Z$ dominance (Z -Higgsino coupling $\sim v_u \lambda_u (\Lambda_u)$ or $\sim v_d \lambda_d (\Lambda_d)$)


Correlation: $g - 2, \mu \rightarrow e\gamma, \mu \rightarrow e$



Dipole A_2 dominates, correlation strong, $g - 2$ large

Summary

- MRSSM: enlarged Higgs and Gauge sector
- λ_d, Λ_d enhancement for $g-2$
- Correlation between $g-2$, $\mu \rightarrow e\gamma$ and $\mu \rightarrow e$



- Dipole A_2 dominance
 → strong correlation between the low-energy lepton observables
- Future experiment results:
 $g-2$ (fermilab), $\mu \rightarrow e\gamma$ (MEG-II), $\mu \rightarrow e$ (COMET, Mu2E)