



Searches for BSM Physics in Rare B-decays at CMS

Martino Margoni

Università di Padova and INFN

on behalf of the CMS Collaboration


- Motivation
- $B \rightarrow \mu\mu$
- $B \rightarrow K^{(*)}\mu\mu$
- Constraints on New Physics

Motivation

Rare B decays: New Physics probes

- Search for deviations from Standard Model (SM) predictions due to virtual contributions of new heavy particles in loop processes
 - Compare experimental results with very precise SM expectations (uncertainty usually dominated by QCD)
- The most interesting processes are those that are strongly suppressed in the SM: Leptonic B-decays, FCNC ($K^{(*)} \mu^+ \mu^-$) [but also LFV, CPV in B^0 mixing, c & τ]
 - New Physics (NP) could modify expectations by orders of magnitude [e.g. A. Buras, arXiv:0910.1032]

- Rare B decays can probe high scales potentially sensitive to NP beyond the direct reach of LHC:



$$\Lambda_{\text{NP}} \sim \frac{M_W}{g^2} \sqrt{\frac{16\pi^2}{|V_{ts}^* V_{tb}|}} \sim 10 \text{ TeV}$$

Rare B decays: New Physics probes

Weak decay of hadron M into final state F described via an Effective Hamiltonian expressed by means of Operator Product Expansion:

$$A(M \rightarrow F) = \langle F | H_{eff} | M \rangle = \frac{G_F}{\sqrt{2}} \sum_i V_{CKM}^i C_i(\mu) \langle F | Q_i(\mu) | M \rangle$$

$C_i(\mu)$: Wilson Coefficients (perturbative short distance couplings)

$Q_i(\mu)$: Hadronic Matrix Elements (non-perturbative long distance effects)

NP could modify Wilson Coefficients $C_i(\mu)$ and/or add new operators $Q_i(\mu)$

Complementary information from different rare decays:

$B \rightarrow \mu\mu$: Scalar/Pseudoscalar interactions

$B \rightarrow K^{(*)} \mu\mu$: Vector/axial interactions


$$\mathcal{B} \rightarrow \mu^+ \mu^-$$

“Measurement of the $B_s^0 \rightarrow \mu^+ \mu^-$ branching fraction and
Search for $B^0 \rightarrow \mu^+ \mu^-$ with the CMS Experiment”

[$L = 5 \text{ fb}^{-1}$ ($\sqrt{S}=7 \text{ TeV}$) + 20 fb^{-1} ($\sqrt{S}=8 \text{ TeV}$)]


Phys. Rev. Lett. 111, 101804 (2013)

$B \rightarrow \mu^+ \mu^-$



- FCNC process forbidden at tree level

- Helicity suppressed $\sim (m_\mu / m_B)^2$



- Cabibbo suppressed $|V_{ts(td)}|^2$
 + $BR \sim 10^{-9}$: Probe the SM!


- SM: suppression of B_d over $B_s \sim (|V_{td}|/|V_{ts}|)^2$:

$$BR(B_s \rightarrow \mu^+ \mu^-) = (3.2 \pm 0.2) 10^{-9} \quad [\text{Buras et al., Eur. Phys. J C72, 2172}]$$

$$BR(B_d \rightarrow \mu^+ \mu^-) = (0.11 \pm 0.01) 10^{-9}$$

Uncertainties from f_{B_s} (lattice), $V_{tb} V_{ts}$, m_t , τ_{B_s}

$B \rightarrow \mu^+ \mu^-$



- Golden mode to search for New Physics with scalar/pseudo-scalar interactions

- NP scenarios in the **extended Higgs sector**:

- May enhance or suppress the BR wrt SM
- Show different $\tan \beta$ dependence:
 - MSSM: $BR \sim \tan^6 \beta / M_A^4$
 - 2 Higgs Doublet Models: $BR \sim \tan^4 \beta$

[e.g. M. Ciuchini, Run2 B Physics
CMS Workshop, Venice April 2014]


- $BR(B_d \rightarrow \mu^+ \mu^-)$ vs $BR(B_s \rightarrow \mu^+ \mu^-)$:

Test of **Minimal Flavor Violation**: general structure of SM FCNC is preserved, flavor violation depends only on CKM

$$B \rightarrow \mu^+ \mu^-$$


Signal:

- Two isolated muons from a secondary vertex
- $M(\mu^+ \mu^-) \sim M(B^0_{s(d)})$
- Momentum aligned with flight direction



BKG:


- Combinatorial from uncorrelated B semileptonic decays
- Physical:
 - Peaking $B \rightarrow hh'$ ($h = \text{misidentified } K, \pi$) ($BR \sim 10^{-7}/10^{-5}$)
 - Non Peaking $B \rightarrow h\mu\nu$, $B \rightarrow h\mu\mu$, $\Lambda_b \rightarrow p\mu\nu$



$$B \rightarrow \mu^+ \mu^-$$

Strategy:

- Use of dedicated dimuon trigger path:



Hardware Trigger:

$P_T(\mu) > 3$ GeV (few kHz)

High Level Trigger 2011 (2012):

Central region ($|\eta| < 1.8$):

$P_T(\mu) > 4$ (3) GeV, $P_T(\mu\mu) > 3.9$ (4.9) GeV,

$4.8 < M(\mu\mu) < 6$ GeV

Forward region ($1.8 < |\eta| < 2.2$):

$P_T(\mu) > 4$ GeV, $P_T(\mu\mu) > 7$, Prob(VTX) $> 0.5\%$

- BDT-based muon identification:

- Exploits kinematic quantities, silicon-tracker fit information and combined silicon/muon track fit information

- Misidentification studied on MC/data control samples





$$(K^0 \rightarrow \pi \pi, \Lambda \rightarrow p \pi, D^* \rightarrow D^0 \pi)$$

- $\epsilon(\pi \rightarrow \mu) < 0.13\%$, $\epsilon(K \rightarrow \mu) < 0.22\%$, $\epsilon(p \rightarrow \mu) < 0.15\%$

$B \rightarrow \mu^+ \mu^-$

Strategy:

- Events selected by means of a MVA exploiting kinematic, vertexing and isolation variables



$$B \rightarrow \mu^+ \mu^-$$

● Strategy:

- Events selected by means of a MVA exploiting kinematic, vertexing and isolation variables
- Measure event yields from an unbinned fit to $M(\mu\mu)$
- BR obtained relative to the normalization channel $B^+ \rightarrow K^+ J/\psi$ to avoid systematics from cross section & luminosity, and reduce efficiency uncertainty:

$$B(B_s^0 \rightarrow \mu^+ \mu^-) = \frac{Y_S}{Y_N} \frac{\epsilon_N}{\epsilon_S} \frac{f_U}{f_S} B(B^+ \rightarrow K^+ J/\psi \rightarrow K^+ \mu^+ \mu^-)$$

Y_S, Y_N	Signal and Normalization Yields	$B(B^+) = (6.0 \pm 0.2) 10^{-5}$
ϵ_S, ϵ_N	Signal and Normalization Efficiencies	
$\frac{f_U}{f_S} = 0.256 \pm 0.020$	Ratio between B^+ and B_s^0 fragmentation functions [LHCb, JHEP 04 (2013) 001]	



- Data/MC agreement checked on $B_s \rightarrow J/\psi \phi$ control sample

$B \rightarrow \mu^+ \mu^-$

Strategy:



- Combinatorial BKG from Side Bands extrapolation
- Semileptonic & Peaking BKG estimated normalizing to $B^+ \rightarrow K^+ J/\psi$

$$N(Y \rightarrow X) = \frac{B(Y \rightarrow X)}{B(B^+ \rightarrow K^+ J/\psi)} \frac{\epsilon(X)}{\epsilon(B^+)} \frac{f_Y}{f_U} N(B^+ \rightarrow K^+ J/\psi)$$



- Peaking BKG checked with independent analysis of $B \rightarrow hh$

$B \rightarrow \mu^+ \mu^-$: Results



Results:

$$\text{BR}(B_s \rightarrow \mu^+ \mu^-) = (3.0^{+0.9}_{-0.8} (\text{stat})^{+0.6}_{-0.4} (\text{syst}) 10^{-9}) \quad (4.3 \sigma \text{ significance})$$

$$\text{BR}(B^0 \rightarrow \mu^+ \mu^-) < 1.1 \times 10^{-9} @ 95\% \text{ CL}$$

Systematics from muon misidentification, BR of rare BKG decays ($\Lambda_b \rightarrow p \mu \nu$) and normalization of peaking BKG

Comparison with other experiments

- ATLAS from 4.9 fb^{-1} using a BDT analysis:

$$\text{BR}(B_s \rightarrow \mu^+ \mu^-) < 1.5 \cdot 10^{-8} @ 95\% \text{ CL}$$


- LHCb from 3 fb^{-1} using a BDT analysis:

$$\text{BR}(B_s \rightarrow \mu^+ \mu^-) = 2.9^{+1.1}_{-1.0} \cdot 10^{-9} \text{ (4.0 } \sigma \text{ significance)}$$


$$\text{BR}(B^0 \rightarrow \mu^+ \mu^-) < 7.4 \times 10^{-10} @ 95\% \text{ CL}$$

[ATLAS-CONF-2013-076]

[LHCb: PRL 111, 101805]




SUSY2014, Manchester 21-26 July 2014



M. Margoni Università di Padova & INFN

Comparison with other experiments

History of a long search



$B^0 \rightarrow \mu^+ \mu^-$:
 Let's wait for next
 LHC Runs

$B \rightarrow \mu^+ \mu^-$: Combination

CMS-PAS-BPH-13-007
LHCb-CONF-2013-012


● Preliminary CMS+LHCb combination

✦ Taking into account correlation from f_s/f_u [LHCb, JHEP 04 (2013) 001]


$$\text{BR}(B_s \rightarrow \mu^+ \mu^-) = (2.9 \pm 0.7) \cdot 10^{-9} \quad (>5 \sigma \text{ significance})$$

$$\text{BR}(B^0 \rightarrow \mu^+ \mu^-) = (3.6^{+1.6}_{-1.4}) \cdot 10^{-10} \quad (<3 \sigma \text{ significance})$$

$B_s \rightarrow \mu^+ \mu^-$





$B^0 \rightarrow \mu^+ \mu^-$




What Next on $B \rightarrow \mu^+ \mu^-$?

[CMS PAS FTR-13-022]



Year	L (fb ⁻¹)	No. of B_s^0	No. of B^0	$\delta B/B(B_s^0 \rightarrow \mu^+ \mu^-)$	$\delta B/B(B^0 \rightarrow \mu^+ \mu^-)$	B^0 sign.	$\delta \frac{B(B^0 \rightarrow \mu^+ \mu^-)}{B(B_s^0 \rightarrow \mu^+ \mu^-)}$
now	20	16.5	2.0	35%	>100%	0.0–1.5 σ	>100%
2018	100	144	18	15%	66%	0.5–2.4 σ	71%
2021	300	433	54	12%	45%	1.3–3.3 σ	47%
2023	3000	2096	256	12%	18%	5.4–7.6 σ	21%

- Expected number of events assuming SM BRs
- High Luminosity-LHC: Inner tracker with improved granularity & muon detector with extended coverage




“Angular analysis and branching fraction measurement of the decay $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ ”

[$L = 5.2 \text{ fb}^{-1}$ ($\sqrt{S} = 7 \text{ TeV}$)]

Phys. Lett. B 727, (2013) 77-100

$$B \rightarrow K^* \mu^+ \mu^-$$




FCNC process forbidden
at tree level, $BR \sim 10^{-6}$: Probe the SM!

- Sensitive to the effects of NP in photon, vector and axial-vector couplings which can enter at the same order as SM contributions

- Complementary information to $B \rightarrow \mu^+ \mu^-$

- Amplitudes expressed using OPE in terms of:

- Hadronic Form Factors (accuracy $\sim 20\%$)
[A. Barucha et al. arXiv 1004.3249]
- Wilson coefficients C_7^{eff} , C_9^{eff} , C_{10}^{eff}
[A. Ali et al., PRD 61 074024, Z. Phys. C 67 417]
- Clean theoretical predictions especially at low $q^2 = m^2(\mu^+ \mu^-)$
- Experimentally clean signature



• Differential Amplitude:

$$\frac{1}{\Gamma} \frac{d^3 \Gamma}{d \cos \theta_K d \cos \theta_l dq^2}$$

$$= \frac{9}{16} \left\{ \left[\frac{2}{3} F_S + \frac{4}{3} A_S \cos \theta_K \right] (1 - \cos^2 \theta_l) \right.$$

$$+ (1 - F_S) \left[2 F_L \cos^2 \theta_K (1 - \cos^2 \theta_l) \right.$$


$$+ \frac{1}{2} (1 - F_L) (1 - \cos^2 \theta_K) (1 + \cos^2 \theta_l) \left. \right.$$

$$\left. + \frac{4}{3} A_{FB} (1 - \cos^2 \theta_K) \cos \theta_l \right\}.$$

- Kinematics of the decay $B \rightarrow V \mu^+ \mu^-$ ($V=K^*$, ϕ , ρ) determined by three angles:
 - + θ_l, θ_K, ϕ
- Event Yields reconstructed in bins of $q^2 = m^2(\mu^+ \mu^-)$

• Observables Include:

- + Differential Branching Ratio dB/dq^2
- + A_{FB} (forward-backward muon asymmetry)
- + F_L (fraction of longitudinally polarized K^*)



• Differential Amplitude:

$$\frac{1}{\Gamma} \frac{d^3 \Gamma}{d \cos \theta_K d \cos \theta_l dq^2}$$

$$= \frac{9}{16} \left\{ \left[\frac{2}{3} F_S + \frac{4}{3} A_S \cos \theta_K \right] (1 - \cos^2 \theta_l) \right.$$

$$+ (1 - F_S) \left[2 F_L \cos^2 \theta_K (1 - \cos^2 \theta_l) \right.$$

$$+ \frac{1}{2} (1 - F_L) (1 - \cos^2 \theta_K) (1 + \cos^2 \theta_l) \left. \right.$$

$$\left. + \frac{4}{3} A_{FB} (1 - \cos^2 \theta_K) \cos \theta_l \right\}.$$

- Kinematics of the decay $B \rightarrow V \mu^+ \mu^-$ ($V=K^*$, φ, ρ) determined by three angles:
 - + θ_l, θ_K, ϕ
- Event Yields reconstructed in bins of $q^2 = m^2(\mu^+ \mu^-)$

- F_S : Fraction of spinless $K\pi$ (S-wave) combination
- A_S : Interference amplitude between S-wave and P-wave decays
 - + $F_S = 0.01 \pm 0.01, A_S = -0.10 \pm 0.01$ fitted on the $B^0 \rightarrow K^* J/\psi$ control sample₂₁



• Strategy:

- Measure event yield Y_S , A_{FB} and F_L from an unbinned simultaneous fit to $M(K\pi\mu\mu)$, $\cos(\theta_K)$ and $\cos(\theta_l)$ in bins of q^2

$$PDF(M, \cos\theta_K, \cos\theta_l) = Y_S \cdot S(M) \cdot S(\cos\theta_K, \cos\theta_l) \cdot \epsilon(\cos\theta_K, \cos\theta_l) \quad \text{Signal}$$

$$+ Y_{Bc} \cdot B_C(M) \cdot B_C(\cos\theta_K) \cdot B_C(\cos\theta_l) \quad \text{Combinatorial}$$

$$+ Y_{Bp} \cdot B_P(M) \cdot B_P(\cos\theta_K) \cdot B_P(\cos\theta_l) \quad \text{Peaking BKG from } B^0 \rightarrow K^* J/\psi(\psi')$$

Y_S, Y_{Bc}, Y_{Bp} Event Yields
 $S(\cos\theta_K, \cos\theta_l), \epsilon(\cos\theta_K, \cos\theta_l)$ Signal 2D angular shape and efficiency
 $S(M), B_C(M), B_P(M)$ Mass PDFs
 $B_C(\cos\theta_{K(l)}), B_P(\cos\theta_{K(l)})$ Angular BKG PDFs

- dB/dq^2 obtained relative to the normalization channel $B^0 \rightarrow K^* J/\psi$:




$$\frac{dB(B^0 \rightarrow K^{*0} \mu^+ \mu^-)}{dq^2} = \frac{Y_S}{Y_N} \frac{\epsilon_N}{\epsilon_S} B(B^0 \rightarrow K^{*0} J/\psi \rightarrow K^{*0} \mu^+ \mu^-)$$

Y_S, Y_N Signal and Normalization Yields
 ϵ_S, ϵ_N Signal and Normalization Efficiencies



● Strategy:

- Measure event yield Y_S , A_{FB} and F_L from an unbinned simultaneous fit to $M(K\pi\mu\mu)$, $\cos(\theta_K)$ and $\cos(\theta_l)$ in bins of q^2



➤ B flavor tagging from $K\pi$ charge




➤ BKG PDFs:

Combinatorial from MC

Peaking parameterized on MC

$B \rightarrow K^* J/\psi(\psi')$




$B \rightarrow K^* \mu^+ \mu^-$: Results



Results consistent with SM

- \blacklozenge Theoretical and experimental errors comparable
- \blacklozenge Systematics from Peaking BKG mass shape, $\cos(\theta_{l,K})$ BKG shape and S-wave contribution

Comparison with other experiments






● LHCb measures the position of the zero-crossing point theoretically clean [JHEP 1308 131]:

$q_0^2 = 4.9 \pm 0.9 \text{ GeV}^2$

In agreement with the SM

[e.g. M. Beneke et al., Eur. Phys. J C47, 625]

Comparison with other experiments



What Next from CMS?


- Results of 8 TeV data analysis expected soon:
 - ⊕ dB/dq^2 , F_L , A_{FB} , A_{FB} zero crossing-point
 - ⊕ Use of new angular variables with small Form-Factor dependence

Constraints on NP

Constraints from $B \rightarrow \mu^+ \mu^-$

SM4: SM with a 4th generation
 MFV: Flavor Violation governed only by CKM matrix

\oplus $BR(B_d)/BR(B_s)$ extremely sensitive probe of NP




● Result in agreement with SM [0.4 σ for B_s and 1.7 σ for B_d]

● The focus now is on $BR(B_d)$ and on the ratio $BR(B_s)/BR(B_d)$

● LHCb (after upgrade): measure ratio @ 35% with 50 fb^{-1}

LHCb $B \rightarrow K^* \mu^+ \mu^-$: Hint of NP?

Variables free from Form Factor contributions [JHEP 05, 137]




3.7 σ discrepancy in P'_5 in

$4.3 < q^2 < 8.68 \text{ GeV}^2$ [PRL 111, 191801]

Possible interpretation as a NP contribution to Wilson coefficient C_9

Low q^2
 $K^* \mu^+ \mu^-$ bins alone

3.9 σ
discrepancy
wrt SM



Combined analysis from $K^* \mu \mu$,

$B \rightarrow X_s \gamma, X_s \mu \mu, K^* \gamma, B_s \rightarrow \mu \mu$


[S. Descotes-Genon et al., PRD 88, 074002]

Difficult to explain with SUSY

Consistent with a Z' with $m \sim 7 \text{ TeV}$

LHCb $B \rightarrow K^* \mu^+ \mu^-$: Hint of NP?

Variables free from Form Factor contributions [JHEP 05, 137]



3.7 σ discrepancy in P'_5 in

$4.3 < q^2 < 8.68 \text{ GeV}^2$ [PRL 111, 191801]

Possible interpretation as a NP contribution to Wilson coefficient C_9

Resulting C_9^{NP} would imply an inclusive $\text{BR}(B \rightarrow X_s l^+ l^-)$ suppression of $\sim 25\%$ in $1 < q^2 < 6 \text{ GeV}^2$ and $q^2 > 14.4 \text{ GeV}^2$

Recent BaBar $\text{BR}(B \rightarrow X_s l^+ l^-)$ result in the high- q^2 region shows a $\sim 2 \sigma$ excess wrt SM prediction in both the $X_s \mu^+ \mu^-$ and $X_s e^+ e^-$ channels

➤ LHCb effect not confirmed by BaBar [PRL 112, 211802]

Conclusions




Conclusions

- Rare B decays are an excellent laboratory for the search for physics beyond the SM
- In the last few years several new measurements from LHC & B-Factories experiments released with impressive experimental precision
- Almost all the results are in agreement with expectations **but some tension is present in some sectors (i.e. $B \rightarrow K^* \mu\mu$, $\tau\nu$, $D^{(*)}\tau\nu, \dots$)**
- **Strong constraints on NP models from flavor measurements**

- Rich program of measurements is expected from LHC/Belle II experiments in the coming years
 - **Chances to discover/understand NP in flavor sector in the near Future?**

Backup

$B \rightarrow K^* \mu^+ \mu^-$: Results




Perturbative region $q_0^2 = 1 < q^2 < 6 \text{ GeV}^2$:

Experiment	F_L	A_{FB}	$d\mathcal{B}/dq^2$ (10^{-8} GeV^{-2})
CMS	$0.68 \pm 0.10 \pm 0.02$	$-0.07 \pm 0.12 \pm 0.01$	$4.4 \pm 0.6 \pm 0.4$
LHCb	$0.65^{+0.08}_{-0.07} \pm 0.03$	$-0.17 \pm 0.06 \pm 0.01$	$3.4 \pm 0.3^{+0.4}_{-0.5}$
BaBar	-	-	$4.1^{+1.1}_{-1.0} \pm 0.1$
CDF	$0.69^{+0.19}_{-0.21} \pm 0.08$	$0.29^{+0.20}_{-0.23} \pm 0.07$	$3.2 \pm 1.1 \pm 0.3$
Belle	$0.67 \pm 0.23 \pm 0.05$	$0.26^{+0.27}_{-0.32} \pm 0.07$	$3.0^{+0.9}_{-0.8} \pm 0.2$
SM	$0.74^{+0.06}_{-0.07}$	-0.05 ± 0.03	$4.9^{+1.0}_{-1.1}$

$B \rightarrow K^* \mu^+ \mu^-$: Hint of NP?

Study additional variables free from FF contributions [JHEP 05, 137 (2013)]




3.7 σ discrepancy in P'_5 in

$4.3 < q^2 < 8.68 \text{ GeV}^2$

Possible interpretation as a NP contribution to Wilson coeff. C_9

Analysis of the full 3 fb^{-1} statistics in progress



P'_5 tension correlated with other minor tensions (too small BRs)



Difficult to explain with SUSY

Consistent with a Z' with $m \sim 7 \text{ TeV}$

Measure other $B \rightarrow K \mu \mu$ decays

$B \rightarrow K \mu^+ \mu^-$

$K^* \mu^+ \mu^-$ tension motivates studies of A_I, A_{FB}



Update of previous A_I measurement (4.4σ discrepancy integrated on q^2)

is now in agreement with SM


- Improvements in efficiency ratio $\varepsilon(K\mu\mu)/\varepsilon(J/\psi K)$
- $\mathcal{B}(J/\psi K^0)/\mathcal{B}(J/\psi K^+)$ correction applied vs K momentum

A_{FB} in agreement with SM

$B \rightarrow K \mu^+ \mu^-$

[LHCb-Paper-2014-006]

● $K^* \mu^+ \mu^-$ tension motivates studies of differential BRs




Decay mode	Measurement	Prediction
$B^+ \rightarrow K^+ \mu^+ \mu^-$	$8.5 \pm 0.3 \pm 0.4$	10.7 ± 1.2
$B^0 \rightarrow K^0 \mu^+ \mu^-$	$6.7 \pm 1.1 \pm 0.4$	9.8 ± 1.0
$B^+ \rightarrow K^{*+} \mu^+ \mu^-$	$15.8 \begin{smallmatrix} +3.2 \\ -2.9 \end{smallmatrix} \pm 1.1$	26.8 ± 3.6

● All the results are “consistent” with SM at $<2.2 \sigma$

● But all of them are lower than the predictions...

What Next on $K^{(*)} \mu^+ \mu^-$ & friends?

- Measurements of related $b \rightarrow d \mu \mu$ channels are welcome to reveal information on Minimal Flavor Violation nature of New Physics




LHCb: $BR(B^+ \rightarrow \pi^+ \mu^+ \mu^-) = (2.4 \pm 0.6 \pm 0.2) 10^{-8}$ in agreement with MFV

[JHEP 12, 125 (2012)]

- $BR(B^+ \rightarrow \pi^+ \mu^+ \mu^-) / BR(B^+ \rightarrow K^+ \mu^+ \mu^-)$ would provide a comparison of $|V_{td}| / |V_{ts}|$ from penguin processes and box processes ($\Delta m_s / \Delta m_d$)
- Improve theoretically very clean measurements of semi-inclusive $B \rightarrow X_{s/d} \mu \mu$, ($X_{s/d} \gamma$)

BaBar $B \rightarrow X_s \ell^+ \ell^-$ [PRL 112, 211802]



● Perturbative region $q_0^2 = 1 < q^2 < 6 \text{ GeV}^2$:


$$BR = (1.60^{+0.41}_{-0.39} \quad +0.17_{-0.13} \quad \pm 0.18) 10^{-6}$$

In agreement with SM $(1.59 \pm 0.11) 10^{-6}$

● q^2 region above $\psi(2S)$:

$$BR = (0.57^{+0.16}_{-0.15} \quad +0.03_{-0.02} \quad \pm 0.0) 10^{-6}$$

$\sim 2 \sigma$ above SM $(0.24 \pm 0.07) 10^{-6}$



✦ LHCb effect not confirmed by BaBar